

## WEST Search History

DATE: Wednesday, January 28, 2004

| Hide?  | Set Name | Query                                     | Hit Count |
|--|----------|---|-----------|
| <i>DB=USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR</i>      |          |   |           |
| <input type="checkbox"/>                                 | L35      | L34 and (gm2 or gm adj 2)                 | 0         |
| <input type="checkbox"/>                                 | L34      | L33 and ganglioside\$                     | 0         |
| <input type="checkbox"/>                                 | L33      | 5891432.pn.                               | 2         |
| <input type="checkbox"/>                                 | L32      | (allogeneic same cancer adj cell) and HSP | 5         |
| <input type="checkbox"/>                                 | L31      | (allogeneic same cancer adj call) and HSP | 0         |
| <input type="checkbox"/>                                 | L30      | L29                                       | 35        |
| <i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR</i> |          |   |           |
| <input type="checkbox"/>                                 | L29      | (allogeneic same cancer) and HSP          | 59        |
| <input type="checkbox"/>                                 | L28      | (allogeneic adj cancer) and HSP           | 0         |
| <input type="checkbox"/>                                 | L27      | (allogenic adj cancer) and HSP            | 0         |
| <input type="checkbox"/>                                 | L26      | (allogenic adj cancer adj cell) and HSP   | 0         |
| <input type="checkbox"/>                                 | L25      | L23 and (gmcsf or gm adj csf)             | 5         |
| <input type="checkbox"/>                                 | L24      | L23 and (gm2 or gm adj 2)                 | 1         |
| <input type="checkbox"/>                                 | L23      | cancer adj cell same hsp90                | 37        |
| <input type="checkbox"/>                                 | L22      | L21 and ganglioside                       | 0         |
| <input type="checkbox"/>                                 | L21      | 6517837 .pn.                              | 2         |
| <input type="checkbox"/>                                 | L20      | L19 and ganglioside                       | 0         |
| <input type="checkbox"/>                                 | L19      | 6576756.pn.                               | 2         |
| <input type="checkbox"/>                                 | L18      | L13 and ganglioside                       | 1         |
| <input type="checkbox"/>                                 | L17      | L15 and (hsp)                             | 0         |
| <input type="checkbox"/>                                 | L16      | L15 and (hsp adj 90 or hsp90)             | 0         |
| <input type="checkbox"/>                                 | L15      | L13 and (gmcsf or gm adj csf)             | 6         |
| <input type="checkbox"/>                                 | L14      | L13 same (gmcsf or gm adj csf)            | 0         |
| <input type="checkbox"/>                                 | L13      | allogeneic adj cancer                     | 13        |
| <i>DB=USPT; PLUR=YES; OP=OR</i>                          |          |   |           |
| <input type="checkbox"/>                                 | L12      | L11                                       | 0         |
| <input type="checkbox"/>                                 | L11      | L9 and radiation                          | 0         |
| <input type="checkbox"/>                                 | L10      | L9 and bcg                                | 0         |
| <input type="checkbox"/>                                 | L9       | L8 and (hsp90 or hsp adj 90)              | 1         |
| <input type="checkbox"/>                                 | L8       | 5981706.pn.                               | 1         |
| <input type="checkbox"/>                                 | L7       | L3 and (hsp adj 90 or hsp90)              | 1         |

*DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR*

☐ L6 L5 1

*DB=PGPB,USPT; PLUR=YES; OP=OR*

☐ L5 L3 same (hsp adj 90 or hsp90) 1

☐ L4 l3 same hsp-90 1


*DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR*

☐ L3 L2 same (gm adj csf or gmcsf) 2

☐ L2 ganglioside same (cancer adj cell) 140

☐ L1 ganglioside same (cancer ad cell) 2112

END OF SEARCH HISTORY



Entrez PubMed Nucleotide Protein Genome Structure PMC Journals Books

Search PubMed for [ ] Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Cancer Biochem Biophys. 1998 Nov; 16(4): 313-32.[Related Articles, Links](#)

Entrez PubMed

[Overview](#)  
[Help | FAQ](#)  
[Tutorial](#)  
[New/Noteworthy](#)  
[E-Utilities](#)

PubMed Services

[Journals Database](#)  
[MeSH Database](#)  
[Single Citation Matcher](#)  
[Batch Citation Matcher](#)  
[Clinical Queries](#)  
[LinkOut](#)  
[Cubby](#)

Related Resources

[Order Documents](#)  
[NLM Gateway](#)  
[TOXNET](#)  
[Consumer Health](#)  
[Clinical Alerts](#)  
[ClinicalTrials.gov](#)  
[PubMed Central](#)[Privacy Policy](#)

## Chemical, metabolic and immunological characterization of gangliosides of human glioma cells.

Maeda Y, Yamaki T, Yoshikawa J, Tatewaki K, Piao H, Yu H, Ibayashi Y, Hashi K.

Department of Neurosurgery, Sapporo Medical University School of Medicine, Japan.


The patterns of ganglioside profiles were studied in 10 human glioma and one melanoma cell lines. Ganglio-series gangliosides, GM3 (NeuAc alpha2-3Gal beta1-4Glc beta1-Cer) and GM2 (GalNAc beta 1-4 (NeuAc alpha2-3)Gal beta1-4Glc beta 1-1Cer), and a neolacto-series ganglioside, sialylparagloboside (SPG) (NeuAc alpha 2-3Gal beta1-4GlcNAc beta1-3Gal beta1-4Glc beta1-1Cer), were the predominant constituents. The activities of the two key enzymes, GM3 synthetase and lactotriaosyl ceramide (Lc3Cer) synthetase, alone did not account for the ganglioside profile. Metabolic labeling with the use of [3H]glucosamine-HCl showed more pronounced difference in the synthetic rate of each ganglioside type, in which GM2 was the most strongly labeled in 7 out of the 10 glioma cell lines. On quantifying the chemical content of GM3 and GM2, the GM3/GM2 molar ratio of above 2.0 was arbitrarily classified into GM3 dominant type (KG-1C and Mewo); the ratio below 0.5 was designated as GM2 dominant type (H4, U138MG, U373MG, T98G and A172); and the ratio between 0.5 and 2.0 was regarded as GM3 and GM2-co-dominant type (U87MG, Hs683, SW1088 and U118MG). Subsequently, the capabilities of the antibody binding to these gangliosides were examined in native forms in the cell membrane and in chemically-isolated forms. The intensity of reaction against chemically isolated GM3 and GM2 gangliosides was dependent on the quantity, and GM2 was more reactive than GM3; however, the reactivities on the cell surface did not correlate with the chemical content indicating other factors to influence their immunoreactivities.

PMID: 9925280 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI | NLM | NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act | Disclaimer](#)

Jan 20 2004 07:12:30



Entrez PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Books

Search PubMed for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources

Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

☐ 1: Biochim Biophys Acta. 1990 Aug 6; 1045(3): 239-44.

Related Articles, Links

## Gangliosides in the human glioma cell line U-118 MG grown in culture or as xenografts in nude rats.

Fredman P, Mansson JE, Bigner SH, Wikstrand CJ, Bigner DD, Svennerholm L.

Department of Psychiatry and Neurochemistry, Gothenburg University, St. Jorgen Hospital, Hisings Backa, Sweden.

This study was undertaken to characterize gangliosides in the human glioma cell line U-118 MG. The cell line was grown both in cell culture and as xenografts in nude rats. A common finding in both culture and xenograft cells was the high proportion of the lactoseries ganglioside 3'-LM1, approximately one third of the total ganglioside sialic acid. Otherwise, there were marked differences between the two cell sources. The cells grown in culture had a more simple ganglioside pattern than those grown in xenografts. In the latter instance, more complex gangliosides of the lactoseries, including 3'8'-LD1, sialyllactonorhexaosylceramide and a branched structure with two terminal NeuAc alpha 2-3Gal beta 1-4GlcNAc chains, and the gangliotetraose series were found. Another marked difference involved GM2, which in the cultured cells was a major fraction, indicating that the synthesis of the gangliotetraose series gangliosides in the former stopped at the level of GM2. These results show that the ganglioside composition of a glioma cell line is strongly influenced by environmental factors.

PMID: 2386796 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk  
NCBI | NLM | NIH  
Department of Health & Human Services  
Freedom of Information Act | Disclaimer

Jan 20 2004 07:12:30